

Geochronology and Tectonic Setting of Volcanic-Hosted Massive Sulphide Deposits in Mainland SE Asia

Khin Zaw^{*a} Mohd Basril Iswadi Basori^b Sebastien Meffre^a R.R. Large^a Charles Makoundi^a, Clive Burrett^c, Hai Thanh Tran^d, Chun Kit Lai^e

^aCODES Centre of Ore Deposits and Earth Sciences, University of Tasmania, Hobart, Australia,

^bGeology Program, Faculty of Science and Technology, National University of Malaysia (UKM),

^cPalaeontological Research and Education Centre, Mahasarakham University, Mahasarakham, Thailand

^dDepartment of Geology, Hanoi University of Mining and Geology, Hanoi, Viet Nam

^eUniversiti Brunei Darussalam, Brunei Darussalam

* Corresponding author: Khin.Zaw@utas.edu.au

ABSTRACT

Volcanic-hosted massive sulphide (VHMS) deposits are formed during the rifting of oceanic ridges or back-arc basins. The SE Asia region is characterised by an assembly of crustal plates or microcontinents that rifted off from northern margin of Gondwana during the Phanerozoic. These microplates or terranes drifted northwards on the Tethyan Oceans and accreted or collided with the Eurasian margin. Throughout the long history of Gondwana supercontinent break-up and subsequent accretion of these crustal terranes, amalgamation and assembly phases of mainland SE Asia, various rifting, subduction, opening and closure of backarc basins, ophiolitic obduction and arc-continent/continent-continent collisions have occurred [1]. The associated complex and multiple tectonomagmatic-metamorphic-hydrothermal interactions among the SE Asia continental fragments have generated several VHMS deposits (e.g., Duc Bo deposit in Vietnam, Tasik Chini deposit in central Malaysia and Leymyetna and Mahar San and Bawdwin deposit in Myanmar and Wetar in Indonesia) [1,2]. The Cu-Pb-Zn Duc Bo deposit is located about 10km from the Coast and 9km south of Tam Ky, the Capital City of Quang Nam Province in central Vietnam. The deposit is hosted in Ordovician-Silurian mafic-felsic volcanic and metavolcanic rocks such as metadacite and metabasalt, and strongly deformed and metamorphosed to Amphibolite facies. The host rocks yielded LAICP-MS zircon age of 417.3 ± 7 Ma to 459.5 ± 9.7 Ma. The Tasik Chini district in central Malaysia has a long history of iron-manganese-barite mining and hosts the Bukit Botol and Bukit Ketaya VHMS ore lenses [3,4]. These ore lenses were formed within weakly deformed felsic-dominated Permian volcanics that are part of a mixed volcano-sedimentary rock succession. The stringer and massive sulphide generally define the lower mineralized zones, whereas the barite, Fe-Mn layers define the stratigraphically upper mineralized zones. The main sulphide phases are pyrite, chalcopyrite, sphalerite, rare galena, and trace Sn-, Au-, and Ag-bearing minerals, with the latter two confined to the massive sulphide and barite zones. LAICP-MS U-Pb zircon dating of rhyolites from the Bukit Botol deposit yields Early Permian (286 ± 4 to 292 ± 3 Ma) ages. Similarly, the zircon U-Pb age results for the Bukit Ketaya rhyolites reveal Early Permian (286 ± 2 to 288 ± 4 Ma) ages [3]. The Bawdwin deposit in northern Myanmar is the largest producing Pb-Zn-Ag VHMS deposit in SE Asia and occurs in Cambro-Ordovician volcanoclastic pile, Numtu Township, northeastern Myanmar and the deposit is relatively undeformed and least metamorphosed. Formation of these VHMS deposits in SE Asia is related to the protracted history and development of Paleo- to Meso-Tethyan Oceans and there is a significant potential for future discovery of these deposit types in SE Asia. Further research is required to improve understanding of the basic geology (particularly to establish time-space relations of magmatic, tectonic and hydrothermal alteration history) for better targeting of the potentials and prospective grounds in the region for mineral exploration.

References

[1] Khin Zaw et al. (2014). *Gondwana Research* 26, 5–30. [2] Khin Zaw (2017). Chapter 24 In: Barber, A.J., Khin Zaw & Crow, M.J. (eds) *Myanmar: Geology, Resources and Tectonics*. Geological Society, London, *Memoirs*, 48, 531–556. [3] Mohd Basril Iswadi Basori et al. (2016). *International Geology Review*, 58, 929–948. [4] Mohd Basril Iswadi Basori et al. (2019). *Ore Geology Reviews*, <https://doi.org/10.1016/j.oregeorev.2019.102947>.

Keywords: VHMS, Permian, Cambrian, Tectonics, Geochronology, Lead-Zinc-copper-silver