
Some Experimental Results of the Station Cone Algorithm Compared to the Simplex Algorithm for Linear Programming

C. N. Nguyen^{1*} and H. T. Le²

DOI: 10.9734/bpi/tpmcs/v10/7037D

Abstract

In this paper we introduce a new variant of station cone algorithm to solve linear programming problems. It uses a series of interior points O_k to determine the entering variables. The number of these interior points is finite and they move toward the optimal point. At each step, the calculation of new vertex is a simplex pivot. The proposed algorithm will be a polynomial time algorithm if the number of points O_k is limited by a polynomial function. The second objective of this paper is to carry out experimental calculations and compare with simplex methods and dual simplex method. The results show that the number of pivots of the station cone algorithm is less than 30 to 50 times that of the dual algorithm. And with the number of variables n and the number of constraints m increasing, the number of pivots of the dual algorithm is growing much faster than the number of pivots of the station cone algorithm. This conclusion is drawn from the computational experiments with $n \leq 500$ and $m \leq 2000$. In particular we also test for cases where $n = 2, m = 100\,000$ and $n = 3, m = 200\,000$. For case where $n = 2$ and $m = 100\,000$, station cone algorithm is given no more than 16 pivots. In case of $n = 3, m = 200\,000$, station cone algorithm has 23 pivots. The test has confirmed the trend that as the number of variables and constraints increases, the number of pivots of the simplex algorithm increases more rapidly than the number of pivots of the station cone algorithm.

Keywords: Linear programming; simplex method; dual problem; station cone.

1 Introduction

Linear programming (LP) is considered as one of the greatest inventions of mathematics in the 20th century. And there are two mathematicians who are regarded as the founders of the LP: Soviet mathematician Leonid Kantorovich (19 January 1912 – 7 April 1986) and American mathematician George Dantzig (November 8, 1914 – May 13, 2005).

In 1939, for the first time, Leonid Kantorovich studied the problem of planning production. And he came up with a mathematical model approach. He set up the mathematical model for the production planning problem along with the solution. The Kantorovich work - "Mathematical methods of organizing and planning production" [1] is recorded as the original appearance of linear programming.

But the important milestone of linear programming as a new field of mathematics was in 1947, when George Dantzig introduced the simplex algorithm. After its discovery by Dantzig in 1947 [2] the simplex method was unrivaled, until the late 1980s, for its utility in solving practical linear programming problems. The computational experiments show that the simplex method is efficient in practice [3,4,2,5]. Nevertheless, there exists a class of linear programming problems for which the simplex method takes an exponential number of steps [6].

In 1979 [7] Khachiyan introduced the ellipsoid method which run in polynomial time (a bound of $O(n^5L)$ arithmetic operations on number with $O(nL)$ digits). Khachiyan's algorithm was of landmark importance for

¹Hanoi Institute of Mathematics, 18 Hoang Quoc Viet, Hanoi, Vietnam.

²Faculty of Informatics Technology, University of Mining and Geology, Hanoi, Vietnam.

*Corresponding author: E-mail: dr.chu.vga@gmail.com;

establishing the polynomial time solvability of linear programs. Despite its major theoretical advance, the ellipsoid method had little practical impact as the simplex method is more efficient for many classes of linear programming problems [8,9].

In 1984 [10] Karmarkar proposed a new projective method for linear programming problems which not only improved Khachiyan's theoretical worst-case polynomial bound but in fact promised dramatically practical performance improvement over simplex method. Karmarkar's algorithm falls within the class of interior point methods. In contrast to the simplex method, which finds the optimal solution among the vertices of the feasible set, the interior point method moves through the interior of the feasible region and reaches the optimal solution only asymptotically. Stimulated by Karmarkar's algorithm a variety of interior point methods were developed for linear programming [11,12].

There are several important open problems in the theory of linear programming, the solution of which would represent fundamental breakthrough in mathematics. In the recent survey on linear programming [13] M.J. Todd has mentioned unsolved problems: Is there a polynomial pivot rule for the simplex method? The immense efficiency of the simplex method in practice, despite its exponential time theoretical performance, hints that there may be variations of simplex algorithm that run in polynomial time.

Therefore, we set ourselves the following 3 purposes: The first purpose is to search the new algorithm more efficiently than the simplex algorithm; The second purpose is to find the polynomial pivot rule for the variation of simplex algorithm; The third purpose is to conduct experimental calculations to compare the newly found algorithm with the simplex algorithm.

In this paper, we present an algorithm, which can be considered a variant of the dual simplex method. In the next section, we introduce the station cone concept, which plays a key role in our algorithm. How to select the leaving variable is presented in the section 3. In section 4, we show how to choose the entering variable - this is an important key to the efficiency of the algorithm. Section 5 devoted to algorithm description. The result of experimental calculation is presented in section 6. A few comments are given in section 7.

2 Station Cone

Consider a linear programming problem in the matrix form

$$\begin{aligned} & \text{Max } \langle c, x \rangle \\ & x \in P := \{x | Ax \leq b, x \geq 0\}, \end{aligned} \tag{2.1}$$

where $c \in \mathbb{R}^n, A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, \forall x \in \mathbb{R}^n$. Let A_1, A_2, \dots, A_m denote the row vectors. Through this paper we suppose that (2.1) and its dual problem are non-degenerated. We also suggest the feasible region P of (2.1) has strict interior points. For simplicity of argument, we assume that the matrix A has full column rank n and $n < m$.

Let $I_n = \{i_1, i_2, \dots, i_n\} \subset \{1, 2, \dots, m\}$ such that the vectors $A_i, i \in I_n$ are linear independent. This means the vector $A_i, i \in I_n$ establish a basis of \mathbb{R}^n . Therefore any vector $A_l \in \mathbb{R}^n$ can be expressed as a linear combination of the vectors $A_{i_k}, i_k \in I_n$. Let λ_{i_k} be the linear coefficient of the vector A_l in the basis $A_{i_k}, i_k \in I_n$, then

$$a_{lj} = \sum_{k=1}^n \lambda_{i_k} a_{i_k j}, \quad j = 1, 2, \dots, n, \quad l = 1, 2, \dots, m.$$

Consider the system of homogeneous linear inequalities

$$A_{i_k} x \leq 0, \quad i_k \in I_n. \tag{2.2}$$

Definition 1. *The linear inequality*

$$A_I x \leq 0 \tag{2.3}$$

is called the consequent linear inequality of the system (2.2) if and only if all the solutions of the system (2.2) satisfy the linear inequality (2.3).

We need the following well known result in theory of linear inequalities.

Theorem 2.1. *The linear inequality (2.3) is a consequent linear inequality of the system (2.2) if and only if*

$$A_I = \sum_{k=1}^n \lambda_{i_k} A_{i_k}, \lambda_{i_k} \geq 0, i_k \in I_n$$

Definition of station cone. *Let polyhedral cone M be defined by system of linear inequalities*

$$\begin{aligned} A_{i_1} x &\leq b_{i_1}, \\ A_{i_2} x &\leq b_{i_2}, \\ &\dots \\ A_{i_n} x &\leq b_{i_n}, \end{aligned}$$

where $A_{i_1}, A_{i_2}, \dots, A_{i_n}$ are linear independent. Then M is called a station cone if the vector c is a nonnegative linear combination of the vectors $A_{i_1}, A_{i_2}, \dots, A_{i_n}$. Then the vertex \bar{x} of the station cone M is called a station solution and the vectors $A_{i_1}, A_{i_2}, \dots, A_{i_n}$ is called a basis of a station cone.

Therefore, geometrically it can be seen that all the station cones lie on one side of the objective function $\langle c, x \rangle$ at their vertices (see Fig 1: M^1, M^2, M^3, M^4, M^5 are station cones and M^6, M^7, M^8, M^9 are not station cones). In other words, the solutions of the system of linear inequalities that create the station cones satisfy the inequality $\langle c, x \rangle \leq \langle c, x^* \rangle$, whereas x^* is the vertex of the station cones. This is equal to the fact that the inequality $\langle c, x \rangle \leq \langle c, x^* \rangle$ is the consequent inequality of the system of the linear inequalities, which formulate the station cone. This also means that the vector c is the nonnegative linear combination of the basic vectors of the station cone.




Fig. 1. Station and non-station cones

Theorem 2.2. *If the station solution \bar{x} satisfies all the constraints of the problem (2.1) then \bar{x} is an optimal solution.*

3 Leaving Variable

Let $A_{i_1}, A_{i_2}, \dots, A_{i_n}$ be the basis of the station cone and

$$c = \sum_{k=1}^n \lambda_{k0} A_{i_k},$$

$$A_j = \sum_{k=1}^n \lambda_{kj} A_{i_k}, \quad j = 1, 2, \dots, m.$$

Then from definition 2.1 follows that: $\lambda_{k0} \geq 0, \quad \forall k = 1, 2, \dots, n$.

From now on we assume that all λ_{k0} are strictly positive, i.e. $\lambda_{k0} > 0, \quad k = 1, 2, \dots, n$.

It is obvious that $\lambda_{k0} > 0, \quad k = 1, 2, \dots, n; \quad \lambda_{k0} = 0, \quad k = n+1, \dots, m$ is a basis solution of the dual problem of (2.1):

$$\begin{aligned} &\text{Min } \langle b, \lambda \rangle \\ &A^T \lambda \geq c^T \\ &\lambda \geq 0, \end{aligned} \tag{3.1}$$

where $\lambda \in R^m$. The assumption $\lambda_{k0} > 0, \quad k = 1, 2, \dots, n$ means that the dual problem (3.1) is non-degenerated.

Remark 1. *The vertex of the station cone is a basic solution of the dual problem.*

4 Pendulum Principle and Entering Variable

We find that, if we connect the vertices of the cones to the center of a circle, the vertices will oscillate around the optimal point according to the pendulum principle. Then finally stop at the optimal point. That is one of the main ideas of the station cone algorithm. In other words, the pendulum principle is one of the spinal ideas, from which the station cone algorithm is formed.




Fig. 2. Pendulum principle

Let us approximate the equator of the earth by a polygon with the edge of 1 meter long. Then this polygon has 40 millions edges and 40 millions vertices. Suppose we have to find the maximum of a linear function $cx_1 + cx_2$ over this polygon.

On Fig. 1, let A denote an optimal point, B^1 denote the starting point. Suppose the distance between B^1 and A is 5 million meters. Then the simplex method will produce an optimal solution after 5 million iterations.

Let M^1 be a station cone defined by 2 constraints containing points B^1 and D^1 , where D^1 is on the other side of A with a distance, for examples, 4 million meters to A (see Fig. 2).

We denote by x^1 the vertex of M^1 . Since M^1 is a station cone, it is clear that $cx^1 \geq cx, \forall x \in M^1$. The station cone M^1 will be our starting cone. Starting our algorithm with the operation of connecting x^1 with O , where O is the center of the equator. The segment $[x^1, O]$ will intersect with the boundary of P at B^2 . Replacing the constraint containing B^1 by the constraint containing B^2 we have a new cone M^2 . Repeat the above procedure with M^2 and we have M^3 , etc. (see Fig. 1). The replacement of one constraint by another has to follow the restriction that the new generating cone is a station cone. We note that at each iteration, the distance between two points B^k and D^k defined by two edges of the station cone M^k is reduced by approximately 2 times in comparison with the previous iteration. Therefore the number of the iterations T can be estimated by the following bound

$$T \approx \log_2 \frac{m}{2} \tag{4.1}$$

For our example with $m = 40$ million the formula (4.1) gives

$$T \approx \log_2 \frac{m}{2} = \log_2 2 \cdot 10^7 < 25.$$

The above example shows that our algorithm can produce an optimal solution after around 25 iterations.

4.1 Initial Station Cone

We now proceed to find an initial station cone. We can find an initial station cone M by solving the following system

$$\begin{aligned} A^T \lambda &= c^T, \\ \lambda &\geq 0, \end{aligned} \tag{4.2}$$

where $\lambda \in \mathbb{R}^m$. We can suppose $c^T \geq 0$ because, if some coefficient of c^T is negative then we multiply both sides of the corresponding equation with -1. To find a solution of (4.2), we solve the following big - M problem

$$\begin{aligned} \text{Min} \{ &M_1 y_1 + M_2 y_2 + \dots + M_n y_n \} \\ &A^T \lambda + E y = c^T, \\ &\lambda \geq 0, y \geq 0, \end{aligned} \tag{4.3}$$

Where, $\lambda \in \mathbb{R}^m, y \in \mathbb{R}^n$ and E is the unit matrix of $(n \times n)$ and M_1, M_2, \dots, M_n are significantly large positive numbers. The problem (4.3) has an optimal solution $\lambda^* \geq 0, y^* = 0$. and λ^* is a solution of (4.2).

We also assume that a strict interior feasible solution O of (2.1) is available. If such an initial point is not available then we modify the problem using the usual big – M augmentation [14] as follows:

$$\begin{aligned} \text{Max } & \{ \langle c, x \rangle - Mx_{n+1} \} \\ & Ax - ex_{n+1} \leq b, \\ & x, x_{n+1} \geq 0. \end{aligned} \tag{4.4}$$

Where $e = (1, 1, \dots, 1)^T \in \mathbb{R}^m$ and M is a significantly large positive number.

Let $x_{n+1}^0 > \max \{ 0, -b_1, -b_2, \dots, -b_m \}$. Then $(0, \dots, 0, x_{n+1}^0)^T$ is a strict interior feasible solution of (4.4) which is in the same form as (2.1).

4.2 Initial interior point

Let O be a strict interior point of P . Denoted by $O^i, i = 1, 2, \dots, n$ the projections of O onto n facets of the station cone M^k . Let $H_i, i = 1, 2, \dots, n$ be the intersection points of the boundary of

P and the segments $O, O^i, i = 1, 2, \dots, n$. Then the new point O^* will be calculated by the following formula

$$O^* = \frac{1}{n+1} \left(\sum_{i=1}^n H_i + O \right) \tag{4.5}$$

5 Station Cone Algorithm

5.1 Initialization

Determine the starting station cone M . Calculate the point O^* by formula (4.5).

Let $M^k = M; O = O^*$.

5.2 Step ($k = 1, 2, \dots$)

If the vertex x^k of the station cone M^k is a feasible point of P , then x^k is an optimal solution. In the contrary case, select the inequality $A_s x \leq b_s$ for entering the station cone and define the inequality $A_r x \leq b_r$ for leaving the station cone. Determine the new station cone $M^{\{k+1\}}$ with the vertex $x^{\{k+1\}}$. Go to next step $k = k + 1$.

Remark 2. *Except for the calculation for finding the entering variable, each step of algorithm 1 is a simplex pivot.*

With the assumption that the dual problem (3.1) of (2.1) is non-degenerated, we hence have the following

Theorem 2.6.

The above algorithm produces an optimal solution after a finite number of iterations.

Proof. Follows from the theorems 2.3, 2.4, 2.5.

6 Computational Experiences

The above proposed station cone algorithm has been tested, using MatLab, on a set of randomly generated linear problems [15] of the form

$$\begin{cases} \text{Max } \langle c, x \rangle \\ Ax \leq b, \end{cases} \quad (6.1)$$

where $c = (1, 1, \dots, 1) \in R^n$, A is the full matrix of $(n \times m)$ with a_{ij} is randomly generated from the interval $[0, 1)$, the vector b has been chosen such that the hyperplanes $\langle A_i, x \rangle = b_i$, $i = 1, \dots, m$ are tangent to the sphere $(0, 1)$ with center at origin and radius $r = 1$. To ensure that (6.1) has a finite optimal solution we add the constraints

$$x_i \leq 1, \quad i = 1, 2, \dots, n. \quad (6.2)$$

The optimal solution and objective function value of ((6.1) - (6.2)) have been retested by simplex algorithm from MatLab.

Function **Data01.m** randomly generates the input data for the problems and stores the matrix A and, vector b in the data base form **Dat01.mat**. Function **Alg01.m** solves the problem by a new proposed algorithm1 and function **Simplex01.m** itself is the simplex algorithm from the optimization toolbox of MatLab.

Test results are shown in the tables below (SCA: Station Cone Algorithm).

Table 1. $n = 2, 3$ and $500 \leq m \leq 100000$

n	m	Problem	Pivots		Ratio (SIMPLEX/SCA)
			SIMPLEX	SCA	
2	500	1	257	9	28.5
	1000	1	518	8	64.8
	2000	1	1000	10	100
	3000	1	1540	11	140
	5000	1	2505	13	192.6
	10000	1	4955	14	353.9
	20000	1	9967	14	711.9
	50000	1	25043	15	1669.5
	100000	1	50314	16	3144.6
3	500	1	44	12	3.6
	1000	1	60	15	4
	2000	1	98	13	7.5
	3000	1	104	18	5.7
	5000	1	149	18	8.2
	10000	1	174	18	9.7
	20000	1	284	17	16.7
	50000	1	423	21	20.1
	100000	1	626	22	28.5
	150000	1	779	18	43.2
	200000	1	912	23	39.7

Table 2. $150 \leq n \leq 300, 200 \leq m \leq 700$

<i>n</i>	<i>m</i>	Problem	Pivots		Ratio (SIMPLEX/SCA)
			SIMPLEX	SCA	
150	200	1	13282	1385	8.230
		2	10385	1531	
		3	11493	1357	
		Average	11720	1424	
150	250	1	12834	1710	7.291
		2	13714	1950	
		3	12672	1720	
		Average	13073	1793	
200	300	1	26367	2628	9.326
		2	24800	2941	
		3	27010	2813	
		Average	26059	2794	
250	300	1	35942	3387	11.047
		2	36978	3434	
		3	40686	3473	
		Average	37869	3473	
250	500	1	66942	5751	11.801
		2	62302	5608	
		3	68747	5422	
		Average	66003	5593	
300	600	1	108448	7964	13.6172
350	700	1	157099	11007	14.2726

Table 3. $n = 300, 400, 500; m = 1000, 2000$

<i>n</i>	<i>m</i>	Problem	Pivots		Ratio
			DUAL SIMPLEX	SCA	DUAL SIMPLEX/SCA
300	1000	1	227 215	8 952	26.44
400	1000	1	388 676	13 266	29.29
500	1000	1	583 464	21 033	27.74
100	2000	1	997 853	21 807	45.75

7 Conclusions

7.1 The above tested examples show that the number of pivots of the station cone algorithm is significantly smaller than the simplex and dual methods.

7.2 The test has confirmed the trend that as the number of variables and constraints increases, the number of pivots of the simplex algorithm increases more rapidly than the number of pivots of the station cone algorithm. Therefore, it is necessary to carry out calculations with larger examples.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Kantorovich LV. Mathematical methods of organizing and planning production; 1939.
- [2] Dantzig GB. Progress in linear programming ORSA Journal on Computing. 1963;6(1):15-22.

- [3] Bixby RE. Implementing the simplex method: The initial basis ORSA Journal on Computing. 1992;4: 267-284
- [4] Bixby RE. Progress in linear programming ORSA Journal on Computing. 1994;6(1):15-22.
- [5] Forrest JJ, Goldfarb D. Steepest-edge simplex algorithms for linear programming Mathematical Programming. 1992;57:341-374.
- [6] Klee V, Minty GJ. How good is the simplex algorithm? Inequalities III ed Shisha O. (Academic Press). 1972;159-175.
- [7] Khachiyan LG. A polynomial algorithm in linear programming (in Russian), Doklady Akademiia Nauk SSSR 224 pp 1093-1096. English translation: Soviet Mathematics Doklady. 1979;20:191-194.
- [8] Balinski ML. Mathematical programming: journal, society, recollections History of Mathematical Programming: a Collection of Personal Reminiscences, ed J.-K. Lenstra, A. H. G. Rinnooy Kan et A. Schrijver (CWI et North-Holland Publishing Company, Amsterdam). 5-18.
- [9] Smale S. On the average number of the simplex method of linear programming. Mathematical Programming. 1983;27:241-262.
- [10] Karmarkar NK. A new polynomial-time algorithm for linear programming Combinatorica. 1984;4:373-395.
- [11] Murty KG. A new practically efficient interior point method for LP algorithmic operations research. 2006;1:3-19.
- [12] Luenberger DG, Ye Y. Linear and Nonlinear Programming (Springer Berlin); 2008.
- [13] Todd MJ. The many facets of linear programming Mathematical Programming. 2002;91:417-436.
- [14] Murty KG. A gravitational interior point method for LP Opsearch. 2005;42(1):28-36.
- [15] Paparrizos K, Samaras N, Stephanides G. An efficient simplex type algorithm for sparse and dense linear programs European Journal of Operational Research. 2003;148:323-334.

Biography of author(s)



Dr. C. N. Nguyen

Hanoi Institute of Mathematics, 18 Hoang Quoc Viet, Hanoi, Vietnam.

He was born on July 07, 1954 in Dien Chau district, Nghe An province. In 1977, he graduated from the Faculty of Cybernetics, Kiev National University. In 1980, he defended his Ph.D thesis in Mathematics - Physics, also at Kiev National University. From 1981-2010, Dr. Chu worked at the Institute of Mathematics, Vietnam Academy of Science and Technology. From 2011 to 2016, he was the Chairman of the Board of Directors of East Asia University of Technology, and also held the position of Vice Rector of East Asia University of Technology. Since 2017, he has been the Chairman of the Board of Directors of SMV Company. In addition, from 2007 to 2011, he was General Secretary of the Vietnam Golf Association. At the present, he is the Vice President of the Hanoi Golf Association. Areas of research include: The integer solutions of the system of linear inequalities; Linear programming; Integer programming; Nonlinear programming. Other areas of interest include: journalism, poetry and song writing. Currently, he is a well-known political writer on social media in Vietnam.



H. T. Le

Faculty of Informatics Technology, University of Mining and Geology, Hanoi, Vietnam.

She was born on September 15, 1964 in Huong Khe district, Ha Tinh province. She graduated from the Mathematics Faculty of Da Lat University in 1985. Her scientific career began in 1991 as a lecturer at the Faculty of Mathematics, Hanoi University of Mining and Geology. In 2009, she successfully defended her PhD thesis in Mathematics - Physics at the Hanoi Institute of Mathematics. In 2010, she moved to the position of Head of Department of Economic Informatics, and Deputy Dean of Faculty of Information Technology of the Hanoi University of Mining and Geology. In 2016, she was awarded the title of Associate Professor by the Ministry of Education and Training of Vietnam. Her field of study is mathematical and information technology methods in economics and geology. The main results obtained include: algorithms to solve linear programming problems, algorithms to find the projection point of any point on a convex set, and algorithms to solve systems of linear equations. In the field of geology, she has obtained a number of important results such as: establishing data recovery algorithms for maps, and building mineral resource management and exploitation programs.

Along with the research, she lectures daily at the Faculty of Information Technology. She is in charge of graduation guidance for students and graduate students, as well as successfully completing the responsibilities of the Head of the Department of Economic Informatics and the responsibility of Deputy Dean of the Faculty of Information Technology of the Hanoi University of Mining and Geology.

© Copyright (2021): Author(s). The licensee is the publisher (B P International).

DISCLAIMER

This chapter is an extended version of the article published by the same author(s) in the following journal.
IOP Conf. Series: Journal of Physics: Conf. Series, 1489: 012016, 2020.